

Standardization of Clinical Decision Making for the Conduct of Credible Clinical Research in Complicated Medical Environments

Alan H. Morris, M.D., Thomas D. East, Ph.D., C Jane Wallace, BSN, Meg Franklin, BSN, Laura Heerman, BSN, Tupper Kinder, BS, Matt Sailor, M.S., Debra Carlson, B.S., Richard Bradshaw, B.S.

Pulmonary Division, LDS Hospital, 8th Avenue and C Street, Salt Lake City, Utah 84143

ABSTRACT

The likelihood that past experience will produce correct guides to current practice depends on the signal-to-noise ratio for the clinical problem of interest. If the signal-to-noise ratio is high, the decision will be sound and patient benefit likely to occur. If the signal-to-noise ratio is low, as is commonly the case with difficult clinical decisions, then personal experience and the best intentions will not assure sound clinical decisions. When the probability of benefit cannot be quantified, clinicians in complex settings are in danger of being misled by data and experience. Quantifiable probabilities established by group experiment or observation will be necessary for clinical decisions that can be expected to confer benefit on the patient. Explicit methods are necessary for interventions that can be replicated in experiments or in practice. Computerized protocols force the articulation of explicit clinical care methods and standardize clinical decision making. We have developed explicit, rule-based protocols, implemented them in our hospital, exported them to other hospitals, and successfully achieved a rigorous experimental environment in the clinical ICU. Exportation of such explicit methods may narrow the gap between efficacy (university hospital) and effectiveness (community hospital) research results.

INTRODUCTION

Medicine is and has traditionally been an empiric undertaking [1-5], based, therefore, on experience, including experimentation. The scientific rigor of the empiric observation is the key concern. Experimental method figures prominently in the evaluation of scientific rigor. The randomized clinical trial (RCT) is widely regarded as the most rigorous clinical investigative method for therapy evaluation [6-14]. The Cardiac Arrhythmia Suppression Trial provides a striking example of the value of an empiric RCT, the results of which reversed standard practice that had been securely based on extensive pathophysiologic understanding [15, 16]. It is only one of many therapies initially enthusiastically embraced and widely applied, only to be abandoned when rigorous studies revealed them to be useless or even harmful [9, 12]. Since clinicians can easily be misled by experience, rigorous evaluations of critical care interventions should be pursued and should be made the basis of practice guidelines.

Reproducibility (replication of results by different investigators) is a key requirement of new information before it becomes incorporated into the

body of accepted scientific knowledge [12, 17, 18]. Replication of scientific work can only take place if the original method is explicit. Without clear articulation of essential methodologic details, it is not possible to reproduce a clinical study. Because of the multiple variables that may influence the outcome of critically ill patients, the challenge of articulating an explicit method in adequate detail is daunting, but it is achievable [19-21]. The demanding task of assuring the consistent implementation of the method by clinicians in the unpredictable clinical setting is even more challenging.

Two major elements, the patient and the clinical caregiver, determine the intensity of patient care and the patient outcome. Both the patient and the clinical caregiver are sources of random noise and of systematic noise (bias). The patient contributes noise because of uncontrollable host factors and because of disease etiology, severity, extent, and duration. Local factors influence the patient's disease and the spectrum of clinical problems. The patient identification and selection process is quite imperfect and may incorporate much local bias due to the prejudices of individual clinicians and investigators. This bias is the result of many factors, among which are characteristics of local clinical environments and failure of the medical community to establish broadly accepted specific definitions of many diseases [22, 23]. The clinical caregiver response to the patient also introduces both random noise and bias. Bias is injected into the response of clinical caregivers because of strongly held opinions based on many factors that influence behavior, including general and local cultural factors, local technical abilities, background, training, and experience. These biases can be important non-experimental co-interventions. They can influence patient outcome and confound the interpretation of clinical trial results leading to inappropriate inferences about the effect of experimental interventions.

The detection of an association between an input signal of interest and an outcome measure requires that the signal of interest be capable of separation from other, unwanted, signals with which it may be confused or by which it may be obscured. A common measure of this capability is the signal-to-noise ratio. Unless the signal-to-noise ratio exceeds 1, the signal will be undetectable. For interventions with large signal-to-noise ratios, uncontrolled observations or small clinical trials can be definitive (e.g., the introduction of penicillin for pneumococcal

pneumonia, or lemon juice for scurvy [24]). However, for clinical with low signal-to-noise ratios, uncontrolled observations may be misleading [25].

The internal validity of clinical trials requires equal treatment of the study groups [9, 27 and 28 in 18, 26]). Non-double-blinded studies must be scrutinized for comparability of the non-investigative co-interventions in the experimental treatment arms. However, many interventional critical care studies, including those that incorporate mechanical ventilation techniques, cannot be double-blinded. In these situations particularly, process control, standardization of clinical decision making, and control of the clinical environment become important in establishing the clinical environment as a clinical laboratory equal to the task of conducting the requisite systematic observations under rigorous experimental conditions. The use of point-of-care decision support tools to standardize clinical decision making (achieve process control) is effective and promising, but not widely available [25, 29, 30].

STANDARDIZED DECISIONS

Since treatments must be applied in a uniform manner to comparable patients before one can evaluate the outcome of a particular medical intervention, the standardization of clinical decision making is of importance [12]. Virtually all clinical trials employ protocols. These protocols include definitions, patient selection criteria, procedural rules, and guidelines for conduct of the trial. They generally provide some specific instructions, but they are not specific enough to adequately control the moment-to-moment process of care. Algorithms usually contain non-specific, judgment-requiring suggestions like "optimize PEEP" or "maximize antibiotic therapy. Clinical algorithm texts and other published guidelines also contain many such general instructions [31-35]. While general instructions provide guidelines and are of value for their conceptual content, they are not executable and fail to standardize clinical decision making. The application of general guidelines is associated with great variation of practice by different clinicians, due to variation in individual clinical practice styles.

Control, measurement, and analysis are the cardinal features of rigorous clinical investigation [36]. Of the three elements, it is control that is usually lacking in the clinical environment with the result that unnecessary variation in clinical care characterizes most critical care clinical trial research [37]. This may result in decreased signal-to-noise ratio for clinical outcomes [25, 38]. The lack of control of clinical care processes may explain much of our failure to define the impact of ICU therapies.

Traditional laboratory research is well known for its control. The misperception that laboratory level control is impossible to achieve in the clinical ICU

environment is widespread. This misperception itself is an important determinant of the clinical research programs mounted in the medical community. It constitutes a paradigmatic view that limits the possibilities for clinical research [4, 5]. The traditional expert (authoritarian) clinical decision making paradigm is based on the incorporation of a large number of current variable values. These are assessed with published information and past physician experience and adjusted for the individual patient [39]. Humans are limited in ability to deal with extensive information. Since many clinical therapeutic decisions made in the complex ICU environment are strongly influenced by items recalled from memory, it is appropriate to question both the suitability of this traditional decision making process and the belief in its superiority over more consistent actuarial (data and rule based) decision techniques [29, 40-58]. A serious reassessment of the widespread clinical belief in the impossibility of *control* in the ICU for both clinical and research processes is needed.

Computerized protocols are an effective way to eliminate unnecessary variation in clinical care and thus impose control on the clinical care process [29]. This control can be expected to reduce noise introduced by the clinical caregiver and increase the signal-to-noise ratio for ultimate clinical outcomes [20, 21, 39, 59]. Humans require support for the persistent commitment to detail and to decision making logic (rules) necessary to generate consistent and reproducible clinical decisions.

PROTOCOL RESULTS

Protocols developed and refined at the LDS Hospital, using the hospital-wide HELP system [60-62], have been exported to PC (Unix and Quinix) platforms and used by other clinical centers in a randomized clinical trial of Acute Respiratory Distress Syndrome (ARDS) patients (AHCPR: HS06594, T. D. East, Ph.D., Principal Investigator). The outcomes of two groups, one with mechanical ventilation clinical decisions standardized with bedside computerized protocols, and the other with mechanical ventilation decisions made exclusively by clinicians, are being compared. About 100 patients have been randomized in 9 centers. Preliminary data analysis indicates a performance during 24,000 hours of application in these 9 centers that is indistinguishable from that at the LDS Hospital where the protocols were developed and where standardized clinical decision making occurred 95% of the 24 hour day in about 100,000 hours of around-the-clock use in about 250 ARDS patients.

There is a natural concern about possible harmful effects of protocol standardized clinical decision making. Data do not support this concern. Patients with ARDS who are supported with computerized protocols have experienced a higher survival than

expected from historical control data [29]. However, there are no data supporting a causal association between computerized protocol use and patient outcome. In fact, from 1987 to 1991 at the LDS Hospital the same unexpectedly high survival was observed in ARDS patients supported with or without computerized protocols that provide decision support for mechanical ventilation [63], and a higher survival during the same period has been reported from the Specialized Center of Research in ARDS at the University of Washington [64]. Computerized protocol decision support of mechanical ventilation appears feasible, safe, and practical (given an appropriate clinical computer infrastructure), but its impact on patient outcome is only currently being explored. Other randomized clinical trials, using less detailed and manually applied protocols, have demonstrated clearly that protocol guided care favorably affects the outcome of patients with thromboembolic disease [65-67].

SUMMARY

The likelihood that past experience will produce correct guides to current practice depends on the signal-to-noise ratio for the clinical problem of interest. If the signal-to-noise ratio is high, the decision will be sound (the strength of inference will be high) and patient benefit likely occur. If the signal-to-noise ratio is low, as is commonly the case with difficult clinical decisions, then the strength of inference will be low and personal experience and the best intentions will not suffice. The quantifiable probabilities established by randomized clinical trials will be necessary for clinical decisions that can be expected to confer benefit on the patient. When the probability of benefit cannot be quantified, clinicians in complex settings are in danger of being misled by data and experience. Computerized protocols standardize clinical decision making. They force the articulation of explicit clinical care methods.

We have developed explicit, rule-based protocols, implemented them in our hospital, exported them to other hospitals, and successfully achieved a rigorous experimental environment in the clinical ICU. Exportation of such explicit methods may narrow the gap between efficacy and effectiveness research results.

REFERENCE

1. Hippocrates. The Art. London: William Heinemann, 1923; vol II;III:192).
2. Aristotle. Metaphysics, Book 1. In: Bambrough R, ed. The Philosophy of Aristotle. New York: A Mentor Book (New American Library), 1963: 1-431.
3. Foss L, Rothenberg K. The second medical revolution. Boston: New Science Library, Shambhala, 1988
4. Hoyningen-Huene P. Reconstructing Scientific Revolutions: Thomas S. Kuhn's philosophy of science. Chicago: The University of Chicago Press, 1993
5. Kuhn T. The structure of scientific revolutions. (2 ed.) Chicago: The University of Chicago Press, 1970(Neurch O, ed. International encyclopedia of unified science; vol 2, Number 2).
6. Chalmers I, Silverman W. Professional and public double standards on clinical experimentation. Contr Clin Trials 1987;8(3):388-391.
7. Chalmers I. Minimizing harm and maximizing benefit during innovation in health care: Controlled or uncontrolled experimentation? BIRTH 1986;13(3):155-164.
8. Donahoe M, Rogers R. An anecdote is an anecdote. . . But a clinical trial is data. Am J Respir Crit Care Med 1994;149:293-294.
9. Hulley S, Cummings S. Designing Clinical Research. Baltimore: Williams and Wilkins, 1988
10. Ilstrup D. Randomized clinical trials: Potential cost savings due to the identification of ineffective medical therapies. Mayo Clin Proc 1995;70:707-710.
11. Peto R, Collins R, Gray R. Large-scale randomized evidence: Large, simple trials and overviews of trials. In: Warren K, Mosteller F, ed. Doing more good than harm: The evaluation of health care interventions. New York: The New York Academy of Sciences, 1993: 314-340. vol 703).
12. Pocock SJ. Clinical Trials: A Practical Approach. New York, NY: John Wiley & Sons, 1983:266.
13. Sandercock P. Collaborative worldwide overviews of randomized trials. In: Warren K, Mosteller F, ed. Doing more good than harm: The evaluation of health care interventions. New York: The New York Academy of Sciences, 1993: 149-155. vol 703).
14. Guntupalli K, Fromm RJ. Clinical trials in lung disease. In: Parker M, Shapiro M, Porembka D, ed. Critical care: State of the art. Anaheim, California: Society of Critical Care Medicine, 1995: 187-205. vol 15).
15. The Cardiac Arrhythmia Suppression Trial (CAST) Investigators. Preliminary report: effect of encainide and flecainide on mortality in a randomized trial of arrhythmia suppression after myocardial infarction. N Engl J Med 1989;321:406-412.
16. The Cardiac Arrhythmia Suppression Trial II Investigators. Effect of the antiarrhythmic agent moricizine on survival after myocardial infarction. N Engl J Med 1992;327(4):227-233.
17. Verdaguer A, Patak A, Sancho JJ, Sierra C, Sanz F. Validation of the medical expert system PNEUMON-IA. Comput Biomed Res 1992;25(6):511-26.

18. Campbell D, Stanley J. Experimental and quasi-experimental designs for research (reprinted from Handbook of Research on Teaching, 1963). Houghton Mifflin Co., 1966:84.
19. East TD, Böhm SH, Wallace CJ, Clemmer TP, Weaver LK, Orme JF, Jr, Morris AH. A successful computerized protocol for clinical management of pressure control inverse ratio ventilation in ARDS patients. *Chest* 1992;101:697-710.
20. East T, Morris A, Wallace C, Clemmer T, Orme JJ, Weaver L, Henderson S, Sittig D. A strategy for development of computerized critical care decision support systems. *Int J Clin Monit Comput* 1992;8:263-269.
21. Henderson S, Crapo R, Wallace C, East T, Morris A, Gardner R. Performance of computerized protocols for the management of arterial oxygenation in an intensive care unit. *Int J Clin Monit Comput* 1992;8:271-280.
22. Lind J. A Treatise of the Scurvy. Edinburgh: Souds Murray and Cochran, 1753:115-117, 148-151, 207, 209.
23. Morris A. ARDS clinical trial issues. In: Haslett C, Evans T, ed. Adult respiratory distress syndrome. London: Chapman & Hall, In Press.
24. Petty T, Bone R, Gee M, Hudson L, Hyers T. Contemporary clinical trials in acute respiratory distress syndrome. *Chest* 1992;101:550-552.
25. Bernard G, Artigas A, Brigham K, Carlet J, Falke K, Hudson L, Lamy M, LeGall J, Morris A, Spragg R. The American-European consensus conference on ARDS: Definitions, mechanisms, relevant outcomes and clinical trial coordination. *Am J Respir Crit Care Med* 1994;149(3):818-824.
26. Guyatt G, Sackett D, Cook D. User's guide to the medical literature: II. How to use an article about therapy or prevention; A. Are the results of the study valid? *JAMA* 1993;270(21):2598-2601.
27. Fisher R. Statistical methods for research workers. London: Oliver & Boyd, 1925.
28. McCall W. How to experiment in education. New York: Macmillan, 1923.
29. Morris A, Wallace C, Menlove R, Clemmer T, Orme JJ, Weaver L, Dean N, Thomas F, East T, Suchyta M, Beck E, Bombino M, Sittig D, Böhm S, Hoffmann B, Becks H, Pace N, Butler S, Pearl J, Rasmussen B. Randomized clinical trial of pressure-controlled inverse ratio ventilation and extracorporeal CO₂ removal for ARDS [published erratum appears in *Am J Respir Crit Care Med* 1994 Mar;149(3 Pt 1):838]. *Am J Respir Crit Care Med* 1994;149(2):295-305.
30. East T, Morris A, Gardner R. Computerized management of mechanical ventilation. In: Grenvik A, Ayres S, ed. Textbook of critical care. 3 ed. Philadelphia, PA: W. B. Saunders Company, 1995: 895-911.
31. Don H, ed. Decision making in critical care. Philadelphia: BC Decker Inc., 1985: Clinical decision making Series.
32. Karlinsky J, Lau J, Goldstein R. Decision making in pulmonary medicine. Philadelphia: BC Decker, 1991.
33. Armstrong R, Bullen C, Cohen S, Singer M, Webb A. Critical care algorithms. New York, NY: Oxford University Press, 1991:100.
34. Boutros AR, Hoyt JL, Boyd WC, Harford CE. Algorithm for management of pulmonary complications in burn patients. *Crit Care Med* 1977;5(2):89.
35. Guidelines-Committee-Society-of-Critical-Care-Medicine. Guidelines for the care of patients with hemodynamic instability associated with sepsis. *Crit Care Med* 1992;20(7):1057-1059.
36. Atkins H. The three pillars of clinical research. *Brit Med J* 1958;(27 December):1547-1553.
37. Thorens J-B, Kaelin R, Joliet P, Chevrolet J-C. Influence of the quality of nursing on the duration of weaning from mechanical ventilation in patients with chronic obstructive pulmonary disease. *Crit Care Med* 1995;23:1807-1815.
38. Morris A. Uncertainty in the management of ARDS: lessons for the evaluation of a new therapy (Editorial). *Intens Care Med* 1994;20(2):87-89.
39. Morris A. Paradigms in management. In: Pinsky M, Dhainaut J, ed. Pathophysiologic foundations of critical care medicine. Baltimore: Williams and Wilkins, 1993: 193-206.
40. Eddy DM. Rationing resources while improving quality: How to get more for less. *JAMA* 1994;272:817-824.
41. Eddy DM. Variations in physician practice, the role of uncertainty. *Health Affairs* 1984;3:74.
42. Eddy DM, Billings J. The quality of medical evidence, implications for quality of care. *Health Affairs* 1988;7:19.
43. Eddy DM. Clinical Decision Making. *JAMA* 1990;263:1265-1275.
44. Eddy DM. Clinical Decision Making: From theory to practice - The challenge. *JAMA* 1990;263:287-290.
45. Eddy DM. Clinical Decision Making: From theory to practice - Anatomy of a decision. *JAMA* 1990;263:441-443.
46. Cannon SR, Gardner RM. Experience with a computerized interactive protocol system using HELP. *Comp Biom Res* 1980;13:399-409.
47. Goldberg L. Simple models or simple processes? Some research on clinical judgments. In: Arkes H, Hammond K, ed. Judgment and decision making: An interdisciplinary reader. Cambridge: Cambridge University Press, 1986: 335-353.
48. Grimm RH, Shimon K, Harlan WR, Estes EH. Evaluation of patient-care protocol use by various providers. *N Engl J Med* 1975;292:507-511.

49. Arkes H. Impediments to accurate clinical judgment and possible ways to minimize their impact. In: Arkes H, Hammond K, ed. *Judgment and decision making: An interdisciplinary reader*. Cambridge: Cambridge University Press, 1986: 582-592.
50. McDonald CJ. Protocol-based computer reminders, the quality of care and the non-perfectability of man. *N Engl J Med* 1976;295:1351-1355.
51. McDonald CJ. Use of a computer to detect and respond to clinical events: Its effect on clinician behavior. *Ann Int Med* 1976;84:162-167.
52. McDonald C, Wilson G, McCabe Jr. G. Physician response to computer reminders. *JAMA* 1980;244:1579-1581.
53. McDonald C, Tierney W. Computer-stored medical records-Their future role in medical practice. *JAMA* 1988; 259:3433-3440.
54. McDonald C. Medical heuristics: The silent adjudicators of clinical practice. *Ann Int Med* 1996;124(1 (Part 1)):56-62.
55. Wirtschafter DD, Scalise M, Henke C, Gams RA. Do information systems improve the quality of clinical research? Results of a randomized trial in a cooperative multi-institutional cancer group. *Comp Biom Res* 1981;14:78-90.
56. Kahneman D, Slovic P, Tversky A. *Judgment under uncertainty: Heuristics and biases*. Cambridge: Cambridge University Press, 1982.
57. Leape L. Error in medicine. *JAMA* 1994;272:1851-1857.
58. Wu A, Folkman S, McPhee S, Lo B. Do house officers learn from their mistakes? *JAMA* 1991;265:2089-2094.
59. Morris A, Gardner R. Computer applications. In: Hall J, Schmidt G, Wood L, ed. *Principles of Critical Care*. New York: McGraw-Hill, 1992: 500-514.
60. Pryor TA, Gardner RM, Clayton PD, Warner HR. The HELP system. *J Med Systems* 1983;7:87-102.
61. Kuperman GJ, Gardner RM, Pryor TA. *HELP: A Dynamic Hospital Information System*. New York: Springer-Verlag, 1991 (Orthner HF, ed. *Computers and Medicine*).
62. Pryor TA. The HELP medical record system. *MD Computing* 1988;5(5):22-33.
63. Suchyta MR, Clemmer TP, Orme JF, Morris AH, Elliott CG. Increased survival of ARDS patients with severe hypoxemia (ECMO criteria). *Chest* 1991; 99:951-955.
64. Milberg J, Davis D, Steinberg K, Hudson L. Improved survival of patients with acute respiratory distress syndrome (ARDS): 1983-1993. *JAMA* 1995;273(4):306-309.
65. Hull RD, Raskob GE, Rosenbloom D. Heparin for 5 Days as Compared with 10 Days in the Initial Treatment of Proximal Venous Thrombosis. *N Engl J Med* 1990;322:1260-1264.
66. Cruikshank MK, Levine MH, Hirsh J, Roberts R, Siguenza M. A Standard Heparin Nomogram for the Management of Heparin Therapy. *Arch Intern Med* 1991;151:333-338.
67. Hull RD, Raskob GE, Rosenbloom D, Lemaire J, Pineo G, Baylis B, Ginsberg J, Panju A, Brill-Edwards P, Brant R. Optimal therapeutic level of heparin therapy in patients with venous thrombosis. *Arch Intern Med* 1992;152:1589-1595.

Acknowledgements: This work was supported by NHLBI grant #HL36787, AHCPR grant #HS06594, the Respiratory Distress Syndrome Foundation and the Deseret Foundation (LDS Hospital).